

# **Ion Enrichment of Snowmelt Runoff Water Caused by Basal Ice Formation**

Gro Lilbæk\* and John W. Pomeroy

Centre for Hydrology, University of Saskatchewan,

117 Science Place, Saskatoon, Saskatchewan, S7N 5C8. \*g.lilbaek@usask.ca

## **ABSTRACT**

Once meltwater reaches the base of a snowpack it can infiltrate the underlying stratum, runoff, or refreeze and form a basal ice layer. Basal ice formation is most common early in melt over saturated or very cold frozen soils. Meltwater becomes enriched in ion concentrations compared to the parent snow due to ion fractionation during thaw and percolation through the snowpack. If ion exclusions occur during basal ice formation, further enrichment of runoff water ion concentrations can occur. The influence of basal ice formation on runoff water chemistry was examined by comparing ion concentrations in runoff water that had sustained basal ice contact, to meltwater before basal ice contact. A series of experiments, involving melting a snowpack in a large insulated box over a cold impermeable substrate in a temperature-controlled room, were carried out. A cooling system at the chemically inert base ensured frozen conditions during snowmelt. Meltwater samples were collected throughout melt from within the snowpack using an extraction tube, and runoff water was collected at the base. Samples of the basal ice were collected after each experiment was completed. All samples were analyzed for major anions and cations. Results showed that formation of basal ice layers enriched the runoff water compared to meltwater before basal ice contact. Ion concentrations in basal ice contact runoff water were up to sixteen times greater than those in no-contact meltwater; however, on average, basal ice contact runoff water

showed 1.5 times the ion concentrations of the no-contact meltwater. The implications are that basal ice formation will act to strongly alter both meltwater ion pathway and concentration. When no basal ice is present, enhanced infiltration causes relatively dilute runoff of only part of the meltwater ion load. When basal ice is present all meltwater runs off and further ion concentration enrichment occurs.

*Keywords:* basal ice; snow chemistry; runoff; frozen ground; enhanced infiltration

## INTRODUCTION

Snowpack metamorphism and refreezing of meltwater within the snowpack relocates incorporated ions to quasi-liquid layers on the grain surface and to veins between grains (Colbeck, 1981). This partitioning, or ion exclusion, occurs because of the individual ions lack of ability to become incorporated into the crystal lattice of ice (Davies *et al.*, 1987). The ability is related to the ion's hydrated radii and capability to form hydrogen bonds. For instance, ions such as  $\text{Na}^+$  and  $\text{Cl}^-$  are incorporated more readily into the crystal lattice than  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$ . These excluded ions can be readily mixed with percolating meltwater, enhancing meltwater ion concentrations.

Studies have shown an initial 2- to 7-fold enrichment in ion concentrations in the meltwater released at the base of the snowpack compared to the bulk composition of the parent snowpack (Johannessen and Henriksen, 1978). A typical ion elution curve shows initial enrichment followed by a rapid decrease until the meltwater is depleted relative to the parent snowpack (e.g. Johannessen and Henriksen, 1978); this is referred to as ion fractionation. Studies have shown that the depletion occurs after approximately one-third of the total melt period. Ion distribution within the parent snowpack, melt-freeze cycles, rain-on-snow events, snow depth, and flow path through the snowpack influence on

fractionation (e.g. Colbeck, 1981; Davies *et al.*, 1982). Consequently, snow meltwater chemistry may be strongly influenced by processes occurring weeks or months prior to melt (Cragin *et al.*, 1996).

Once the meltwater reaches the base of a snowpack it either infiltrates the underlying stratum, runs off, or ponds. Ponding occurs when melt rates exceed infiltration rates; this may result in the formation of a basal ice layer (Woo and Heron, 1981). Basal ice can also form where there is a heat flow from the snow to the underlying stratum, which causes infiltrating water to refreeze. Basal ice layers has been observed in glacial environments, where it is referred to as superimposed ice, and non-glacial environments (e.g. Woo *et al.*, 1982; Marsh and Woo, 1984), but to the knowledge of the authors, no studies have presented the influence that a growing ice layer at the base of the snowpack has on runoff chemistry. Conjectures were made by Marsh and Pomeroy (1999) and Jones and Pomeroy (2001).

Thus, the objective of this paper is to identify whether meltwater flow in contact with a basal ice layer becomes enriched in ion concentrations compared to meltwater, which has no contact with a basal ice layer. It is hypothesized that ion exclusion takes place during basal ice formation and that an enriched brine will be present at the basal ice surface, which mixes with meltwater in contact with it.

## METHODOLOGY

An insulated box, with a cooling system at the base, was used to control snowmelt and basal ice formation (Figure 1). The front of the box had a slit along the base for collection of runoff water, which had sustained basal ice contact (BW). An ABS half-pipe was placed with its base ~30 mm above the slit opening for extraction of meltwater




Figure 1: A) Schematic drawing of the experimental box and its instrumentation (not to scale). B) The actual box placed in the temperature controlled room.

before basal ice contact (SW). The half-pipe covered a tenth of the base area. The base consisted of an inert metal plate, which was underlain by a cooling system that was connected to a temperature controlled refrigerated bath (RTE-DD8, Neslab). The whole setup was tilted 8°.

Variations in snowpack temperature were measured using thermocouples (TC; E- and T-type); they were placed at the base and 10, 20, 30, 40, and 45 cm above the base. A TC (K-type) was used to measure the temperature of the coolant. Averaged temperatures were recorded every 15 minutes by a 23X Campbell Scientific datalogger.

Materials that did not have a chemically inert surface were covered by polyethylene to avoid contamination. All materials used were cleaned thoroughly with deionized distilled water (DDI). Plastic gloves were worn at all times to prevent contamination.

Three experiments were conducted using snow from a site located within city limits (experiment 1) and two sites located in rural areas (experiment 2 and 3). The snow used in experiment 1 and 2 was a few days old, but metamorphism was not pronounced. Fresh snow was used in experiment 3. In all experiments, the snow was composed of loose 1-2

Table 1: *Ion concentrations [ $\mu\text{eq l}^{-1}$ ] in the parent snowpacks.*

Experiment #	H <sup>+</sup>	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	Na <sup>+</sup>	K <sup>+</sup>	Mg <sup>2+</sup>	Ca <sup>2+</sup>
1	0.009	12.7	11.5	5.5	6.2	22.8	73.7
2	0.077	25.3	35.4	15.1	11.0	30.9	84.5
3	0.428	9.8	11.5	3.1	8.4	25.0	13.0

mm crystals. The chemical compositions of the parent snow were dominated by magnesium and calcium (Table 1).

To ensure an initial homogeneous snowpack, the snow was sieved with an 8 mm mesh (HDPE). The depth of the snowpack was 0.45 m, giving a total snow volume of 0.15 m<sup>3</sup> within the box. The densities of the snowpacks were ~350 kg m<sup>-3</sup> in the first experiment, ~410 kg m<sup>-3</sup> in the second, and ~215 kg m<sup>-3</sup> in the third. The snow was then left to reach thermal equilibrium; room temperature was -2 °C. Once thermal equilibrium was reached, the room temperature was increased to +5 °C; initial coolant temperature was set at -5 °C, -4 °C, and -3 °C in each experiment, respectively. Small adjustments of the coolant temperature were made throughout the experiment to ensure formation of basal ice. These adjustments were made based on the TC readings at the base of the snowpack.

During melt, water samples (BW and SW) were collected simultaneously twice a day. Minimum sample size was 60 ml due to analytical methods.

All water samples were stored cold until filtered through 45  $\mu\text{m}$  Nuclepore filter paper. Prior to filtration, pH was measured using an Orion 290A meter with an Orion low maintenance triode. Chloride was measured using ion chromatography (Dionex ICS 2000); detection limit was 0.1 mg l<sup>-1</sup>. Inductively coupled plasma atomic emission spectroscopy (ICP-AES) was used for SO<sub>4</sub><sup>2-</sup> analysis. Detection limit was 0.1 mg l<sup>-1</sup>.

Standard methods for the examination of water and wastewater (21st Ed. 2005. Part 2500 and 3000) were used for  $\text{Cl}^-$  and  $\text{SO}_4^{2-}$  analyses. Analyses of  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ , and  $\text{Ca}^{2+}$  were done with atomic absorption spectroscopy (SpectrAA 220, Varian); detection limits were  $0.1 \text{ mg l}^{-1}$  for all ions. Samples analyzed for cations were acidified with  $\text{H}_2\text{SO}_4$  to pH 2 to limit further chemical reactions. Ion balances were not assessed as analyses of  $\text{NO}_3^-$  and  $\text{NH}_4^+$  could not be done. In-between filtering and analyses, samples were stored at  $-20 \text{ }^\circ\text{C}$ .

## RESULTS

The three experiments differed with respect to snowpack density, melt rate, and temperature at the base of the snowpack. In all of the experiments both density and melt rate increased as melt progressed (Figure 2). The greatest increase in snowpack density was observed in experiment 1 with  $\sim 56 \text{ kg m}^{-3} \text{ d}^{-1}$ . This experiment showed the lowest average melt rate ( $5.9 \text{ mm d}^{-1}$ ) and total change in snow water equivalent, SWE (7.4 mm) of all the experiments. The greatest individual melt rate was observed in experiment 2 with  $17.6 \text{ mm d}^{-1}$ ; average melt rate was  $7.7 \text{ mm d}^{-1}$ . The increase in density was about a fifth of the other experiments;  $\sim 11 \text{ kg m}^{-3} \text{ d}^{-1}$ . The total change in SWE was the greatest in all of the experiments; 25.4 mm. In experiment 3 the rate of increase in density was almost the same as in experiment 1;  $\sim 50 \text{ kg m}^{-3} \text{ d}^{-1}$ . The average melt rate was the highest observed ( $8.2 \text{ mm d}^{-1}$ ) and the total change in SWE was about half that of experiment 2; 13 mm. The average base temperature throughout melt was  $-2 \text{ }^\circ\text{C}$  in experiment 1,  $-0.5 \text{ }^\circ\text{C}$  in exp. 2, and  $-0.2 \text{ }^\circ\text{C}$  in exp. 3.




Figure 2: Variation in snowpack density (A), melt rate (B), and snow water equivalent, SWE (C), during melt.

The timing of when the first water was present for collection varied between experiments. Snowpack structure causes instability of the meltwater wetting front, which results in formation of flow fingers and preferential flow paths within the snowpack (Wankiewicz, 1979). The flow fingers are an important mechanism for rapid delivery of meltwater to the bottom of the snowpack while it is still below  $0\text{ }^{\circ}\text{C}$  (Woo *et al.*, 1982).

In each of the experiments runoff water with basal ice contact (BW) was collected at the base before meltwater with no basal ice contact (SW) was present for collection. The reason being that flow fingers did not intersect with the SW extraction pipe. The first BW was collected after 3 days in experiment 1, 2 days in experiment 2, and 1 day in experiment 3. SW was collected 1-4 days after the initial BW sample; after 1 day in experiment 1, 4 days in experiment 2, and 2 days in experiment 3. Total melt period was shortest for experiment 3 (5 days) followed by experiment 1, which lasted 6 days. Experiment 2 melted over approximately 11 days.

The ion concentrations in the BW and SW were normalized to the corresponding ion concentration in the parent snowpack. The normalized value is the concentration

factor, CF (Johannessen and Henriksen, 1978), and shows enriched conditions when  $CF > 1$  and depleted conditions when  $CF < 1$ .

Examples of the variation in CF during melt for both BW and SW are shown in Figure 3. Only the curves for  $SO_4^{2-}$  and  $Ca^{2+}$  are shown as  $Cl^-$ ,  $Na^+$ ,  $K^+$ , and  $Mg^{2+}$  showed similar trends; just different CFs. General for all experiments were that BW showed more irregular curves than the SW; additionally, the CFs for the BW were predominantly greater than or similar to the SW. The variations in  $H^+$  concentrations were not as distinct as for the other ions, showing similar CFs for both BW and SW throughout melt; for the majority of the melt period CF was less than 1, increasing to just above 1 by the end of the melt.

For most of the ions, the greatest CFs were obtained within the initial third of the melt period. Table 2 shows that in general the greatest CFs were obtained in experiment 1; the lowest values were obtained in experiment 3.

Normalizing the ion concentration in the BW with the corresponding concentration in the SW indicates whether the BW is enriched ( $>1$ ) or depleted ( $<1$ ) compared to the SW. Figure 4 shows the variations in BW:SW ratio throughout melt for each ion. The vertical spread of the points indicates if the ratio changes throughout melt; little spread indicates little variation.

Experiment 1 showed the least variation in ratio throughout melt for all ions. Generally, the BW contained half the  $H^+$  concentration of the SW and at least double the concentration for the rest of the ions. Greater scatter in the ratios were observed in experiment 2, where the ratios initially increased and was followed by a decrease. Only  $SO_4^{2-}$  showed continuous enrichment throughout melt. In the last experiment, the ratios




Figure 3: Fractionation curves showing change in concentration factors,  $CF$ , during melt for the major ions  $SO_4^{2-}$  and  $Ca^{2+}$  in the runoff water (BW, closed circles) and the snowpacks meltwater (SW, open circles). The gray horizontal line represents  $CF=1$ .

Table 2: The greatest concentration factors,  $CF$  [ $(\mu eq\ l^{-1}) (\mu eq\ l^{-1})^{-1}$ ], in the runoff water, BW, and meltwater, SW, for each ion in each experiment.

Experiment #		$CF, H^+$	$CF\ Cl^-$	$CF\ SO_4^{2-}$	$CF\ Na^+$	$CF\ K^+$	$CF\ Mg^{2+}$	$CF\ Ca^{2+}$
1	BW	8.7	19.8	9.8	<b>40.7</b>	1.7	5.5	8.0
	SW	15.7	8.0	3.8	<b>19.5</b>	0.6	2.5	2.8
2	BW	1.9	7.3	8.2	5.7	<b>20.9</b>	5.2	8.2
	SW	1.4	<b>22.2</b>	2.2	2.4	21.8	2.9	3.5
3	BW	1.1	5.1	3.5	<b>9.4</b>	1.6	1.0	5.2
	SW	1.1	4.0	2.7	<b>8.0</b>	2.0	1.7	5.3




Figure 4: Variation in the ion ratio between runoff water, BW, and snow meltwater, SW.

were very close to 1 during the whole melt period. Very little variations were seen for H<sup>+</sup>, Cl<sup>-</sup>, and SO<sub>4</sub><sup>2-</sup>; for the rest of the ions, the ratio increased throughout time.

After termination of the experiment, the residual snow was removed from the box exposing the basal ice layer, BI. The thickness of the BI varied from 80 mm in the first experiment to respectively 40 and 30 mm in the following two experiments.

## DISCUSSION AND CONCLUSION

The influence of a growing basal ice layer on runoff chemistry was examined through three snowmelt experiments differing in snowpack density, melt rate, and basal cooling conditions. Ion concentrations in runoff water that had sustained basal ice contact were compared to ion concentrations in meltwater without basal ice contact. The results showed that contact with a basal ice layer enriched the ion concentrations in the runoff water compared to fractionation solely happening within the snowpack. The implications are that formation of basal ice will alter both flow path and concentration of the ion load released from the snowpack. When no basal ice is present, a non-linear relationship

between meltwater ion load and infiltration rate into frozen soil results in enhanced infiltration, which causes a dilution of the runoff water (Lilbæk and Pomeroy, 2007). When basal ice is present all meltwater runs off (e.g. Jones and Pomeroy, 2001) and further ion concentration enrichment occurs. Thus, in areas with variations in extent of frozen soil great variance may be observed in soil and flow path chemistry.

The experiments exhibited a large variation in the timing of water collection. This was believed to be a result of variations in initial snowpack density, melt rate, flow paths, and basal temperature. Finger flow initially feeds the base of the snowpack with meltwater (e.g. Wankiewicz, 1979), where it refreezes and basal ice is formed. The base temperature influenced the thickness of the growing ice layer. Experiment 1 had the coldest base temperature throughout melt ( $-2\text{ }^{\circ}\text{C}$ ) and grew the thickest basal ice layer (80 mm). The base temperatures were higher in the other two experiments ( $\sim -0.5\text{ }^{\circ}\text{C}$ ) and their basal ice layers were approximately half that of experiment 1 (30-40 mm). These basal ice layers compare to those found in nature (Woo *et al.*, 1982; Marsh and Woo, 1984).

At some point, a state of equilibrium, where refreezing of meltwater releases enough latent heat to increase the temperature of the surrounds to the melting point, is reached and runoff initiates. The combination of the thick basal ice layer, high initial snowpack density, and slow average melt rate in experiment 1 resulted in a later collection of initial runoff water compared to the other experiments. The small variation in timing between experiment 2 and 3 was probably affected by initial snowpack density. The delay between initial runoff water and initial meltwater was believed to be a function

of the arrival of the matrix flow to the lower part of the snowpack and when the equilibrium state was reached for the basal ice.

The degree of enrichment of both runoff water and meltwater compared to the parent snowpack decreased with experiment. The greatest CFs were obtained for all ions, except  $K^+$ , in experiment 1; lowest CFs were obtained in experiment 3. The ion fractionation curves decayed exponentially throughout melt, showing the highest concentrations during the initial part of the melt. Overall, little scatter were observed for the meltwater curves. The runoff water curves, however, showed abrupt increases in enrichment followed by concentrations that were slightly depleted or equal to the meltwater; except in experiment 1. Based on the BW:SW ratio, 84% of the compared samples in experiment 1 showed an enrichment of the runoff water compared to the meltwater; only the ratios for  $H^+$  showed depleted conditions. Average enrichment showed twice the ion concentration in the runoff water compared to the meltwater. In experiment 2, 55% of the samples were enriched and 40% were depleted; average enrichment was  $\sim 1.5$ . The BW:SW ratios in experiment 3 did not show much enrichment as melt proceeded for the anions; only for the cations. Still, 46% of the samples were enriched, showing an average enrichment of  $\sim 1.3$ ; 37% were depleted.

A possible reason for the scatter observed in the runoff water ion concentrations is the flow rate and growth of the basal ice layer. In the first experiment, the basal ice was increasing in thickness throughout melt, causing continuous ion exclusion and thereby enrichment of the runoff water. In experiment 2, a state of equilibrium was reached, during which the meltwater that contacted the base would either refreeze, causing ion exclusion and enriching the surface brine, or become mixed with runoff water causing a

decrease in the enrichment rate. The lack of enrichment in experiment 3 can be explained by a late formation of a basal ice layer so the initial samples just represent the snowpacks meltwater. However, in general, it was found that the surface of the basal ice was enriched in  $\text{Cl}^-$ ,  $\text{K}^+$ , and  $\text{Ca}^{2+}$ , and depleted in  $\text{Na}^+$  and  $\text{Mg}^{2+}$ .

The conducted experiments showed that ion exclusion does take place during basal ice formation, enhancing the enrichment of the runoff water. The degree of enrichment depends on the rate of growth for the basal ice layer.

#### ACKNOWLEDGEMENTS

The authors would like to thank Michael Solohub for helping with the construction of the experimental box used. Also appreciated was analytical help from Barry Goetz and Bobbi Helgason. Funding for the research was provided by Canada Research Chair Program, the Canada Foundation for Innovation, and the Government of Saskatchewan Science and Technology Fund.

#### REFERENCES

- Colbeck, S.C. 1981. A Simulation of the Enrichment of Atmospheric Pollutants in Snow Cover Runoff. *Water Resources Research* **17**(5): 1383-1388.
- Cragin JH, Hewitt AD and Colbeck SC. 1996. Grain-scale mechanisms influencing the elution of ions from snow. *Atmospheric Environment* **30**(1): 119-127.
- Davies TD, Brimblecombe P, Tranter M, Tsiouris S, Vincent CE, Abrahams P and Blackwood IL. 1987. The removal of soluble ions from melting snowpacks. *In: Seasonal snowcovers :physics, chemistry, hydrology*. Ed: Jones HG and Orville-Thomas WJ. D. Reidel Publishing company. Dordrecht ; 337-392.

- Davies TD, Vincent CE and Brimblecombe P. 1982. Preferential Elution of Strong Acids from a Norwegian Ice Cap. *Nature* **300**: 161-163. DOI: 10.1038/300161a0
- Johannessen, M. and Henriksen, A. 1978. Chemistry of snow meltwater: Changes in ion concentration during melting. *Water Resources Research* **14**: 615-619.
- Jones HG and Pomeroy JW. 2001. Early spring snowmelt in a small boreal forest watershed: influence of concrete frost on the hydrology and chemical composition of streamwaters during rain-on-snow events. *Proceedings of the 58th Eastern Snow Conference*: 209-218.
- Lilbæk G and Pomeroy JW. 2007. Modelling enhanced infiltration of snowmelt ions into frozen soil. *Hydrological Processes*. In press.
- Marsh P and Pomeroy JW. 1999. Spatial and temporal variations in snowmelt runoff chemistry, Northwest Territories, Canada. *Water Resources Research* **35**(5): 1559-1567.
- Marsh P and Woo MK. 1984. Wetting Front Advance and Freezing of Meltwater Within a Snow Cover; 1. Observations in the Canadian Arctic. *Water Resources Research* **20**(12): 1853-1864.
- Wankiewicz, A. 1979. A review of water movement in snow. *In: Proceedings: Modeling of snow cover runoff*. Ed: S.C. Colbeck and M. Ray. US Army Cold Regions Research and Engineering Laboratory. 222-252.
- Woo M, Heron R and Marsh P. 1982. Basal Ice in High Arctic Snowpacks. *Arctic and Alpine Research* **14**(3): 251-260.
- Woo MK and Heron R. 1981. Occurrence of Ice Layers at the Base of High Arctic Snowpacks. *Arctic and Alpine Research* **13**(2): 225-230.